

Influenza A Virus, A/Puerto Rico/8-CV8/1934 (H1N1)¹

Catalog No. NR-29035

Product Description: Pooled allantoic fluid from specific pathogen free (SPF) embryonated chicken eggs² infected with influenza A virus, A/Puerto Rico/8-CV8/1934 (H1N1)

Lot³: 61617354

Manufacturing Date: 04APR2013

TEST	SPECIFICATIONS	RESULTS
Identification by Infectivity Using Embryonated Chicken Eggs² Hemagglutination activity using allantoic fluid from infected eggs and 0.5% chicken red blood cells	Positive	Positive
Sequencing of Hemagglutinin Coding Region Hemagglutinin (1046 nucleotides) CV8 mutation confirmation	Consistent with A/Puerto Rico/8-CV8/1934 (H1N1) Confirmed	99% identity with A/Puerto Rico/8-CV8/1934 (H1N1) (GenBank: CY084054) Confirmed
Titer by CEID₅₀ Assay^{4,5} in Embryonated Chicken Eggs²	Report results	1.6 × 10 ⁹ CEID ₅₀ /mL
Sterility (21-day incubation) Harpo's HTYE broth ⁶ , 37°C and 26°C, aerobic Trypticase soy broth, 37°C and 26°C, aerobic Sabouraud broth, 37°C and 26°C, aerobic Blood agar, 37°C, aerobic Blood agar, 37°C, anaerobic Thioglycollate broth, 37°C, anaerobic DMEM with 10% FBS, 37°C and 5% CO ₂	No growth No growth No growth No growth No growth No growth No growth	No growth No growth No growth No growth No growth No growth No growth
Mycoplasma Contamination Agar and broth culture (14-day incubation at 37°C) DNA detection by PCR of extracted Test Article nucleic acid	None detected None detected	None detected None detected

¹Deposited and labeled as influenza A virus, A/Puerto Rico/8-34-CV8/2010 (H1N1)

²10-day-old SPF Embryonated Chicken Eggs acquired from B&E Eggs, York Springs, Pennsylvania

³Grown in the allantoic cavity of embryonated chicken eggs² for 2 days at 35°C in a humidified chamber

⁴The Chicken Embryo Infectious Dose 50% (CEID₅₀) is the dilution of virus that under the conditions of the assay can be expected to infect 50% of the inoculated embryonated chicken eggs, just as a Lethal Dose 50% (LD₅₀) is expected to kill half of the animals exposed. A reciprocal of the dilution required to yield the CEID₅₀ provides a measure of the infectious titer (or infectivity) of a virus preparation.

⁵2 days at 35°C in a humidified chamber

⁶Atlas, Ronald M. *Handbook of Microbiological Media*. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

Date: 11 DEC 2013

Signature:



Title:

Technical Manager, BEI Authentication or designee

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

