

NIH AIDS Reagent Program

20301 Century Boulevard Building 6, Suite 200 Germantown, MD 20874 USA

Phone: 240 686 4740 Fax: 301 515 4015 aidsreagent.org

DATA SHEET

Reagent: Sup-T1 CCR5+ Cells (L23)

Catalog Number: 13784

Lot Number: 200217

С Release Category:

Provided: 1 mL of cells

Post thaw cell count = 8.65×10^6 cells/mL

Post thaw cell viability = 78%

Cell viability increased to 91% after 6 days in culture.

Cell Type: Human T cell lymphoblast

Propagation Medium:

RPMI, 90%; fetal bovine serum, 10%; 3 µg/ml Blasticidin; Glutamine; Penicillin;

Streptomycin

Fetal bovine serum, 90%; DMSO, 10% Freeze Medium:

Morphology: Suspension, Lymphocytic Cell Line

Sterility: Negative for mycoplasma, bacteria, and fungi

Description: SupT1 derived T cell clone engineered to express low levels of CCR5.

Special

SupT1 cells were transduced with CCR5 expressing lentivirus. Transduced cells were Characteristics: sorted, individual clones were selected and expanded in blasticidin media and

characterized for CD4 and CCR5 expression. Sup-T1 CCR5+ Cells (L23) can be used in

HIV infection studies.

These cells were received as part of a group and details about each reagent can be

found in Table 1.

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

REV: 07/17/2020 Page 1 of 2 Table 1: Sup-T1 CCR5+ Cells

Alternate Name: SupT-R5 clone L23

Recommended

Storage:

Keep the reagent in liquid nitrogen.

Contributor: Drs. Anjali Joshi and Himanshu Garg

References:

Joshi, A., Nyakeriga, A. M., Ravi, R. and Garg, H. (2011). HIV ENV glycoprotein-mediated bystander apoptosis depends on expression of the CCR5

co-receptor at the cell surface and ENV fusogenic activity. J Biol Chem, (42), 36404-13.

doi:10.1074/jbc.M111.281659 PUBMED

NOTE:

Acknowledgment for publications should read "The following reagent was obtained through the NIH AIDS Reagent Program, Division of AIDS, NIAID, NIH: Sup-T1 CCR5+ Cells (L23) from Drs. Anjali Joshi and Himanshu Garg (cat# 13784)." Also include the

references cited above in any publications.

Scientists at for-profit institutions or who intend commercial use of this reagent must contact the Office of Research Contracts and Agreements (RCA)

at the following email address: rafael.caraveo@ttuhsc.edu, before the reagent

can be released.

July 17, 2020 **Last Updated**

ALL RECIPIENTS OF THIS MATERIAL MUST COMPLY WITH ALL APPLICABLE BIOLOGICAL, CHEMICAL, AND/OR RADIOCHEMICAL SAFETY STANDARDS INCLUDING SPECIAL PRACTICES, EQUIPMENT, FACILITIES, AND REGULATIONS. NOT FOR USE IN HUMANS.

Page 2 of 2 REV: 07/17/2020