

**Spike Glycoprotein (Stabilized) from SARS-Related Coronavirus 2, B.1.1.7 Variant (9 Mutations) with C-Terminal Histidine Tag, Recombinant from HEK293 Cells**

**Catalog No. NR-55421**

**ACROBiosystems Catalog No. SPN-C52H6**

**Product Description:**

A recombinant form of the spike (S) glycoprotein from severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2), B.1.1.7 variant [United Kingdom (UK) variant; also known as 20B/501Y.V1 or VOC202012/01] was produced by transient transfection in human embryonic kidney HEK293 cells and purified by affinity chromatography. NR-55421 lacks the signal sequence and contains 1195 residues (ectodomain) of the SARS-CoV-2 S glycoprotein; the recombinant protein was modified to remove the polybasic S1/S2 cleavage site (R683A and R685A), stabilized with multiple proline substitutions (F817P, A892P, A899P, A942P, K986P and V987P, wild type numbering) and includes a T4 fibrin trimerization motif and a poly-histidine tag at the C-terminus. NR-55421 is a variant of SARS-CoV-2 which contains the HV69-70del, Y144del, N501Y, A570D, D614G, P681H, T716I, S982A and D1118H mutations in the S glycoprotein as compared to the SARS-CoV-2 reference sequence (GenPept: [QHD43416](#)). NR-55421 lot 4353-211WF1-VR was lyophilized from 103 µL bulk protein in phosphate-buffered saline with 10% trehalose.

**Lot: 4353-211WF1-VR**


**Receipt Date: 13MAY2021**

TEST	SPECIFICATIONS	RESULTS
Appearance	White powder	White powder
Purity SDS-PAGE Analysis	Protein band of interest represents > 90% of total staining intensity	Dominant band of ~ 165 kDa represents > 90% of total staining intensity (Figure 1) <sup>1</sup>
Final Product Amount per vial	Report results	50 µg
Functional Activity by ELISA	Reactive	Reactive <sup>2</sup>
Filtration	0.2 µm sterile-filtered	0.2 µm sterile-filtered
Endotoxin Content (Limulus Amoebocyte Lysate Assay)	< 1.0 EU per µg	< 1.0 EU per µg

<sup>1</sup>The recombinant protein migrated to a slightly larger size than was expected, likely caused by glycosylation common in recombinant spike proteins derived from coronaviruses. For more information, please see Chakraborti, S., et al. "The SARS Coronavirus S Glycoprotein Receptor Binding Domain: Fine Mapping and Functional Characterization." *Virology*, 2 (2005): 73. PubMed: 16122388.

<sup>2</sup>Measured by its binding ability in a functional ELISA. Immobilized NR-55421 at 1 µg/mL can bind human ACE2, Fc tag (ACROBiosystems Cat. No. AC2-H5257) with a linear range of 0.1 to 3 ng/mL.

**Figure 1: SDS-PAGE Analysis**



/Sonia Bjorum Brower/

Sonia Bjorum Brower

06 MAY 2024

Technical Manager or designee, ATCC Federal Solutions

ATCC®, on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected by the contributor to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC®'s knowledge.

ATCC® is a trademark of the American Type Culture Collection.

You are authorized to use this product for research use only. It is not intended for human use.

