

***Enterococcus faecalis* EnGen0303, Strain Ned10**

Catalog No. NR-31987

For research use only. Not for use in humans.

Contributor:

Michael S. Gilmore, Sir William Osler Professor of Ophthalmology, Department of Ophthalmology, Massachusetts Eye and Ear Infirmary, Boston, Massachusetts, USA

Manufacturer:

BEI Resources

Product Description:

Bacteria Classification: *Enterococcaceae*, *Enterococcus*

Species: *Enterococcus faecalis* EnGen0303 (Also referred to as *Enterococcus faecalis*)

Strain: Ned10

Original Source: *Enterococcus faecalis* EnGen0303 (*E. faecalis*), strain Ned10 was isolated in 1961 from a horse in the Netherlands.^{1,2}

Comments: *E. faecalis* EnGen0303, strain Ned10 is reported to be resistant to chloramphenicol and tetracycline.¹ The complete genome of *E. faecalis* EnGen0303, strain Ned10 has been sequenced (GenBank: [AJED00000000](https://www.ncbi.nlm.nih.gov/nuclseq/AJED00000000)).

E. faecalis is a Gram-positive, facultatively anaerobic coccus that is a commensal inhabitant of the gastrointestinal and female genital tract.³ It is also the most frequently isolated species, often as a mono-infection, from root canals of endodontically treated teeth with persistent apical periodontitis.⁴ *E. faecalis* is an opportunistic pathogen and has become a serious concern in hospitals because of its inherent hardiness and high levels of antibiotic resistance.⁵ Virulent strains often express a cytolysin toxin that is encoded on various mobile genetic elements, pathogenicity islands, and conjugative plasmids.⁶

Material Provided:

Each vial contains approximately 0.5 mL of bacterial culture in Tryptic Soy broth supplemented with 10% glycerol.

Note: If homogeneity is required for your intended use, please purify prior to initiating work.

Packaging/Storage:

NR-31987 was packaged aseptically in cryovials. The product is provided frozen and should be stored at -60°C or colder immediately upon arrival. For long-term storage, the vapor phase of a liquid nitrogen freezer is recommended. Freeze-thaw cycles should be avoided.

Growth Conditions:

Media:

Tryptic Soy broth or Brain Heart Infusion broth or equivalent Tryptic Soy agar or Tryptic Soy agar with 5% defibrinated

sheep blood or Brain Heart Infusion agar or equivalent

Incubation:

Temperature: 35 to 37°C

Atmosphere: Aerobic (with or without 5% CO₂) or anaerobic

Propagation:

1. Keep vial frozen until ready for use, then thaw.
2. Transfer the entire thawed aliquot into a single tube of broth.
3. Use several drops of the suspension to inoculate an agar slant and/or plate.
4. Incubate the tube, slant and/or plate for 1 day.

Citation:

Acknowledgment for publications should read "The following reagent was obtained through BEI Resources, NIAID, NIH: *Enterococcus faecalis* EnGen0303, Strain Ned10, NR-31987."

Biosafety Level: 2

Appropriate safety procedures should always be used with this material. Laboratory safety is discussed in the following publication: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, and National Institutes of Health. [Biosafety in Microbiological and Biomedical Laboratories \(BMBL\)](#), 6th ed. Washington, DC: U.S. Government Printing Office, 2020.

Disclaimers:

You are authorized to use this product for research use only. It is not intended for human use.

Use of this product is subject to the terms and conditions of the BEI Resources Material Transfer Agreement (MTA). The MTA is available on our Web site at www.beiresources.org.

While BEI Resources uses reasonable efforts to include accurate and up-to-date information on this product sheet, neither ATCC® nor the U.S. Government makes any warranties or representations as to its accuracy. Citations from scientific literature and patents are provided for informational purposes only. Neither ATCC® nor the U.S. Government warrants that such information has been confirmed to be accurate.

This product is sent with the condition that you are responsible for its safe storage, handling, use and disposal. ATCC® and the U.S. Government are not liable for any damages or injuries arising from receipt and/or use of this product. While reasonable effort is made to ensure authenticity and reliability of materials on deposit, the U.S. Government, ATCC®, their suppliers and contributors to BEI Resources are not liable for damages arising from the misidentification or misrepresentation of products.

Use Restrictions:

This material is distributed for internal research, non-commercial purposes only. This material, its product or its derivatives may not be distributed to third parties. Except as performed under a U.S. Government contract, individuals contemplating commercial use of the material, its products or its derivatives must contact the contributor to determine if a

license is required. U.S. Government contractors may need a license before first commercial sale.

References:

1. McBride, S. M., et al. "Genetic Diversity among *Enterococcus faecalis*." PLoS One 2 (2007): e582. PubMed: 17611618.
2. Gilmore, M.S., Personal Communication.
3. Schleifer, K. H. and R. Kilpper-Bälz. "Transfer of *Streptococcus faecalis* and *Streptococcus faecium* to the Genus *Enterococcus* nom. rev. as *Enterococcus faecalis* comb. nov. and *Enterococcus faecium* comb. nov." Int. J. Syst. Bacteriol. 34 (1984): 31-34.
4. Stevens, R. H., O. D. Porras and A. L. Delisle. "Bacteriophages Induced from Lysogenic Root Canal Isolates of *Enterococcus faecalis*." Oral Microbiol. Immunol. 24 (2009): 278-284. PubMed: 19572888.
5. Arias, C. A. and B. E. Murray. "The Rise of the *Enterococcus*: Beyond Vancomycin Resistance." Nat. Rev. Microbiol. 10 (2012): 266-278. PubMed: 22421879.
6. McBride, S. M., et al. "Genetic Variation and Evolution of the Pathogenicity Island of *Enterococcus faecalis*." J. Bacteriol. 191 (2009): 3392-3402. PubMed: 19270086.

ATCC® is a trademark of the American Type Culture Collection.

