SUPPORTING INFECTIOUS DISEASE RESEARCH

Plasmodium falciparum, Strain 3D7, Clone 10.2

Catalog No. MRA-1271

Product Description:

MRA-1271 is a gametocyte-deficient clonal parasite line derived by targeted disruption of Pfs230 in *Plasmodium falciparum* strain 3D7. MRA-1271 was produced by cultivation of deposited material in fresh human erythrocytes suspended in RPMI 1640 medium supplemented with 10% (v/v) heat-inactivated human serum (pooled Type A), 25 mM HEPES, 2 mM L-glutamine, 2 g/L D-glucose, 27 μ g/mL hypoxanthine and 5 μ g/mL gentamicin. The culture was incubated at 37°C in sealed flasks outgassed with a blood-gas atmosphere (90% N₂, 5% CO₂, 5% O₂) and monitored for parasitemia for 7 days. Every 1 to 3 days, uninfected, leukocyte-filtered, Type O erythrocytes in complete culture medium were added dropwise to the culture as needed and monitored for hematocrit.

Lot: 63449603

Manufacturing Date: 04MAY2015

BEI Resources is committed to ensuring digital accessibility for people with disabilities. This Certificate of Analysis contains complex tables and may not be fully accessible. Please let us know if you encounter accessibility barriers and a fully accessible document will be provided: E-mail: <u>Contact@BEIResources.org</u>. We try to respond to feedback within 24 hours.

TEST	SPECIFICATIONS	RESULTS
Identification by Giemsa Stain Microscopy ¹	Blood-stage parasites present	Blood-stage parasites present
Genotypic Analysis ¹		
Sequencing of Merozoite Surface Protein 2 (MSP2) gene (~ 780 base pairs)	Consistent with P. falciparum	Consistent with <i>P. falciparum</i> (Figure 1)
Phenotypic Analysis		
Gametocytemia (<i>in vitro</i>) ²	Report results	< 0.01%
Pfs230 expression (immunofluorescence assay) ³	Report results	Negative
Level of Parasitemia by Giemsa Stain Microscopy		
Pre-freeze (7 days post-infection) ⁴		
Ring-stage parasitemia	Report results	1.78%
Total parasitemia	≥ 2%	2.96%
Post-freeze (3 days post-infection) ¹		
Ring-stage parasitemia	Report results	1.41%
Total parasitemia	≥ 1%	4.21%
Viability (3 days post-infection) ¹	Growth in infected red blood cells	Growth in infected red blood cells
Sterility (21-day incubation) ¹		
Harpo's HTYE broth, 37°C and 26°C, aerobic⁵	No growth	No growth
Trypticase soy broth, 37°C and 26°C, aerobic	No growth	No growth
Sabouraud broth, 37°C and 26°C, aerobic	No growth	No growth
DMEM with 10% FBS, 37°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, aerobic	No growth	No growth
Sheep blood agar, 37°C, anaerobic	No growth	No growth
Thioglycollate broth, 37°C, anaerobic	No growth	No growth
Mycoplasma Contamination ¹		
DNA detection by PCR	None detected	None detected

¹Testing completed on vialed, post-freeze material.

²Gametocytemia was <0.01% after 18 days of *in vitro* growth, confirming that MRA-1271 should not be classified as a gametocyte-producing strain.
³Using monoclonal anti-*Plasmodium falciparum*, 3D7 230 kDa Gamete Surface Protein (BEI Resources MRA-878A) as primary antibody (1:250) and Goat anti-Mouse IgG (H+L) (Invitrogen™ F-2761) as secondary antibody (1:500).

⁴Testing completed on bulk material prior to vialing and freezing.

⁵Atlas, Ronald M. <u>Handbook of Microbiological Media</u>. 3rd ed. Ed. Lawrence C. Parks. Boca Raton: CRC Press, 2004, p. 798.

b|**e**|**i** resources

Certificate of Analysis for MRA-1271

SUPPORTING INFECTIOUS DISEASE RESEARCH

Figure 1: MRA-1271 MSP2 Sequence

/Sonia Bjorum Brower/ Sonia Bjorum Brower

Technical Manager or designee, ATCC Federal Solutions

ATCC[®], on behalf of BEI Resources, hereby represents and warrants that the material provided under this certificate has been subjected to the tests and procedures specified and that the results described, along with any other data provided in this certificate, are true and accurate to the best of ATCC[®]'s knowledge.

ATCC[®] is a trademark of the American Type Culture Collection. You are authorized to use this product for research use only. It is not intended for human use.



26 OCT 2023